

The static and dynamic polarisability, and the Stark and black-body radiation frequency shifts of the molecular hydrogen ions H_2^+ , HD^+ , and D_2^+

S. Schiller

Institut für Experimentalphysik, Heinrich-Heine-Universität Düsseldorf, 40225 Düsseldorf, Germany

D. Bakalov

*Institute for Nuclear Research and Nuclear Energy,
Tsarigradsko chaussée 72, Sofia 1784, Bulgaria*

A. K. Bekbaev

Al-Farabi Kazakh National University, 050012, Almaty, Kazakhstan

V. I. Korobov

Joint Institute for Nuclear Research, 141980, Dubna, Russia

(Dated:)

We calculate the DC Stark effect for three molecular hydrogen ions in the non-relativistic approximation. The effect is calculated both in dependence on the rovibrational state and in dependence on the hyperfine state. We discuss special cases and approximations. We also calculate the AC polarisabilities for several rovibrational levels, and therefrom evaluate accurately the black-body radiation shift, including the effects of excited electronic states. The results enable the detailed evaluation of certain systematic shifts of the transitions frequencies for the purpose of ultra-high-precision optical, microwave or radio-frequency spectroscopy in ion traps.

I. INTRODUCTION

The molecular hydrogen ions represent a family of simple quantum systems that are amenable both to high-precision ab-initio calculations [1, 2] and to high-precision spectroscopy. Therefore, they are of great interest for the determination of fundamental constants [3], for tests of the time- and gravitational-potential-independence of fundamental constants [4, 5], and for tests of QED [2]. On the experimental side, after early pioneering work on uncooled trapped ions and ions beams [6–8], the sympathetic cooling of trapped molecular hydrogen ions [9, 10] has opened up the window for high-precision radio-frequency, rotational, and rovibrational spectroscopy. Precision infrared laser spectroscopy of two rovibrational transitions has been achieved [3, 11], and the fundamental rotational transition has also been observed [12].

Because of the advances in experimental accuracy, and in order to open perspectives for future work directions, it has become important to evaluate the systematic effects on the transition frequencies. It is an advantage of the molecular hydrogen ion family that the sensitivities to external fields can be calculated ab-initio. The systematic effects treated so far include the Zeeman shift [13–15], the electric quadrupole shift [16], and the black-body radiation (BBR) shift [18]. The electric polarisability of the rovibrational levels of the molecular hydrogen ions has been of interest for a long time. It was computed with high accuracy for a subset of levels by several authors, in particular [19–24]. These calculations used adiabatic or non-adiabatic wave functions. Ref. [20] reviews the experimental and theoretical values for the ground state of H_2^+ and D_2^+ . A particularly accurate calculation of the polarisability of H_2^+ in its ground state was performed by one of the present authors, by including the relativistic corrections [25]. The dependence of the polarisability on the hyperfine state has only recently been obtained [26], for the case of HD^+ , and it was shown that the dependence is very significant. These results have permitted a first analysis of the potential for ultra-high accuracy spectroscopy of HD^+ and its suitability as an optical clock [16, 17].

In the present paper, extensive calculations of the polarisability are presented. Its dependence on the hyperfine state is derived in a more elegant way and discussed in depth, both for HD^+ and H_2^+ , since it is of great relevance for experiments.

While the BBR shift is tiny, it will eventually become of relevance for experiments requiring the highest levels of accuracy, such as the mentioned test of time-independence of fundamental constants. Therefore, this shift is also computed in detail. Results for H_2^+ are presented for the first time. In addition, the case of HD^+ is treated extensively, in view of the current experimental interest in this molecule.

This paper is structured as follows: In Sec. 2 we briefly review the calculation approach for the polarisability of the molecular hydrogen ions, neglecting spin effects. We define the effective Hamiltonian and present the tables of polarisabilities. In Sec. 3 we introduce the hyperfine structure and discuss the computation of the DC Stark shift in dependence of the spin state. We also give a number of useful approximations. Sec. 4 presents detailed results for a

large number of hyperfine states potentially relevant for high-precision spectroscopy. In Sec. 5 we discuss the energy level shifts induced by the oscillating (AC) electric field of the black-body radiation, which we accurately evaluate by taking into account the precise frequency dependence of the polarisability.

II. EVALUATION OF THE POLARISABILITY

A. Non-relativistic polarisability. Spin-independent spatial considerations

For the purposes of evaluating the systematic effects in spectroscopy, it is at present sufficient to use the non-relativistic approximation to the polarisability. Therefore, we start from the non-relativistic Schrödinger equation:

$$(H_0 - E)\Psi_0 = 0, \quad H_0 = -\frac{1}{2M_1}\nabla_1^2 - \frac{1}{2M_2}\nabla_2^2 - \frac{1}{2m_e}\nabla^2 + \frac{1}{R} - \frac{1}{r_1} - \frac{1}{r_2}, \quad (1)$$

where M_1 and M_2 are the masses of the nuclei (proton or deuteron), R is the internuclear distance, r_1 and r_2 are the distances from nuclei 1 and 2 to the electron, respectively. The state $\Psi_0 = |v L\rangle$ is the unperturbed state characterized by the vibrational and rotational quantum numbers v , L , and E_0 is its energy.

The interaction with an external electric field $\mathbf{E}$ in the dipole interaction form is expressed by

$$V_p = -\mathbf{E} \cdot \mathbf{d}, \quad \mathbf{d} = e [Z(\mathbf{R}_1 + \mathbf{R}_2) - \mathbf{r}], \quad (2)$$

where $\mathbf{d}$ is the electric dipole moment of the three-particle-system, $\mathbf{R}_{1,2}$ and $\mathbf{r}$ are the position vectors of the nuclei and of the electron with respect to the center of mass.

Since the static or quasi-static electric fields present in an ion trap, and also the electric field of the radiation from continuous-wave lasers and from the black-body environmental radiation are typically weak, it is sufficient to apply second-order perturbation theory for the calculation of the polarisability. The energy shifts that result are typically at the level of 1 Hz, orders of magnitude smaller than the rotational or hyperfine splittings. For effects of higher-order in the external electric field, see Ref. [24].

The change of energy due to the polarisability of a molecular ion is expressed by

$$\begin{aligned} E_p^{(2)} &= \langle \Psi_0 | V_p (E_0 - H_0)^{-1} V_p | \Psi_0 \rangle \\ &= E^i E^j \langle \Psi_0 | d^i (E_0 - H_0)^{-1} d^j | \Psi_0 \rangle \\ &= -\frac{1}{2} \alpha_d^{ij} E^i E^j, \end{aligned} \quad (3)$$

where α_d^{ij} , the polarisability tensor of rank 2, has been introduced,

$$\alpha_d^{ij} = -2 \langle \Psi_0 | d^i (E_0 - H_0)^{-1} d^j | \Psi_0 \rangle. \quad (4)$$

The static dipole polarisability tensor is then reduced to scalar, α_s , and tensor, α_t , terms, which may be expressed in terms of three contributions corresponding to the possible values of the rotational angular momentum quantum number of the intermediate state: $L' = L \pm 1$, or $L' = L$.

$$\begin{aligned} a_+ &= \frac{2}{2L+1} \sum_p \frac{\langle v L \| \mathbf{d} \| p (L+1) \rangle \langle p (L+1) \| \mathbf{d} \| v L \rangle}{E_0 - E_p}, \\ a_0 &= -\frac{2}{2L+1} \sum_p \frac{\langle v L \| \mathbf{d} \| p L \rangle \langle p L \| \mathbf{d} \| v L \rangle}{E_0 - E_p}, \\ a_- &= \frac{2}{2L+1} \sum_p \frac{\langle v L \| \mathbf{d} \| p (L-1) \rangle \langle p (L-1) \| \mathbf{d} \| v L \rangle}{E_0 - E_p}, \end{aligned} \quad (5)$$

where L is the rotational quantum number of the state under consideration, E_p is the non-relativistic energy of the intermediate state $|p L'\rangle$.

The polarisability tensor may be expressed as

$$\alpha_d^{ij} = \delta_{ij}\alpha_s + \alpha_t \left\langle \Psi_0 | L_i L_j + L_j L_i - \frac{2}{3} \delta_{ij} \mathbf{L}^2 | \Psi_0 \right\rangle, \quad (6)$$

where L_i are the Cartesian components of the rotational angular momentum operator, $\mathbf{L}^2 = L_x^2 + L_y^2 + L_z^2$, and

$$\begin{aligned} \alpha_s &= \frac{1}{3}(a_+ + a_0 + a_-), \\ \alpha_t &= -\frac{a_+}{2(L+1)(2L+3)} + \frac{a_0}{2L(L+1)} - \frac{a_-}{2L(2L-1)}. \end{aligned}$$

We may also define longitudinal ($\alpha_{\parallel}$) and transverse ($\alpha_{\perp}$) polarisabilities

$$\alpha_{\parallel} = \alpha_s + \alpha_t \left\langle \Psi_0 | 2L_z^2 - \frac{2}{3} \mathbf{L}^2 | \Psi_0 \right\rangle \quad (7)$$

$$\alpha_{\perp} = \frac{1}{2}(\alpha_d^{xx} + \alpha_d^{yy}) = \alpha_s + \alpha_t \left\langle \Psi_0 | L_x^2 + L_y^2 - \frac{2}{3} \mathbf{L}^2 | \Psi_0 \right\rangle = \alpha_s - \frac{1}{2} \alpha_t \left\langle \Psi_0 | 2L_z^2 - \frac{2}{3} \mathbf{L}^2 | \Psi_0 \right\rangle. \quad (8)$$

The definition of $\alpha_{\perp}$ as given above is reasonable, since axial symmetry requires that the matrix elements of L_x^2 and of L_y^2 are equal. Thus, the polarisabilities $\alpha_{\parallel}$ and $\alpha_{\perp}$ actually involve the expectation value of only a single operator, which has an alternative representation as the 0-component of the rank-2 tensor $\{\mathbf{L} \otimes \mathbf{L}\}_2$,

$$2L_z^2 - \frac{2}{3} \mathbf{L}^2 = \sqrt{\frac{8}{3}} \{\mathbf{L} \otimes \mathbf{L}\}_{20}. \quad (9)$$

In Sec. III, we will evaluate the polarisabilities of the hyperfine states of a given ro-vibrational level. The approximation we will use consists in introducing the polarisability operator, which holds in a manifold of given L ,

$$\begin{aligned} \hat{\alpha}_d^{ij}(v, L) &= \alpha_s(v, L) + \alpha_t(v, L) \left[L_i L_j + L_j L_i - \frac{2}{3} \delta_{ij} \mathbf{L}^2 \right], \\ \hat{\alpha}_{\parallel}(v, L) &= \alpha_s(v, L) + \alpha_t(v, L) \sqrt{\frac{8}{3}} \{\mathbf{L} \otimes \mathbf{L}\}_{20}, \\ \hat{\alpha}_{\perp}(v, L) &= \alpha_s(v, L) - \frac{1}{2} \alpha_t(v, L) \sqrt{\frac{8}{3}} \{\mathbf{L} \otimes \mathbf{L}\}_{20}. \end{aligned}$$

Here, we have included the explicit dependence of the coefficients α_s , α_t on the vibrational and rotational quantum numbers v , L . In the following, we will explicitly consider the polarisability anisotropy operator,

$$\hat{\alpha}_{\parallel} - \hat{\alpha}_{\perp} = \alpha_t(v, L) \frac{3}{2} \sqrt{\frac{8}{3}} \{\mathbf{L} \otimes \mathbf{L}\}_{20} = 3\alpha_t(v, L) \left(L_z^2 - \frac{1}{3} \mathbf{L}^2 \right). \quad (10)$$

B. Numerical results

Wave functions of the rovibrational states in the molecular hydrogen ions are obtained by using the variational approach expounded in Ref. [25]. Briefly, the wave function for a state with a total orbital angular momentum L and of a total spatial parity $\pi = (-1)^L$ is expanded as follows:

$$\begin{aligned} \Psi_{LM}^{\pi}(\mathbf{R}, \mathbf{r}_1) &= \sum_{l_1+l_2=L} \mathcal{Y}_{LM}^{l_1 l_2}(\hat{\mathbf{R}}, \hat{\mathbf{r}}_1) G_{l_1 l_2}^{L\pi}(R, r_1, r_2), \\ G_{l_1 l_2}^{L\pi}(R, r_1, r_2) &= \sum_{n=1}^N \left\{ C_n \operatorname{Re} [e^{-\alpha_n R - \beta_n r_1 - \gamma_n r_2}] + \right. \\ &\quad \left. D_n \operatorname{Im} [e^{-\alpha_n R - \beta_n r_1 - \gamma_n r_2}] \right\}, \end{aligned} \quad (11)$$

TABLE I: Polarisabilities of the HD^+ molecular ion, in atomic units.

v	$L = 0$		$L = 1$		$L = 2$		$L = 3$		$L = 4$		$L = 5$	
	α_s		α_s	α_t	α_s	α_t	α_s	α_t	α_s	α_t	α_s	α_t
0	395.30633		3.99015	175.48275	4.00956	13.82797	4.03878	3.19075	4.07794	1.10141	4.12721	0.47319
1	462.65271		4.70314	205.20067	4.72694	16.14340	4.76278	3.71557	4.81084	1.27799	4.87136	0.54642
2	540.68636		5.56925	239.58035	5.59871	18.81611	5.64313	4.31921	5.70273	1.48001	5.77786	0.62955
3	631.40288		6.63284	279.47585	6.66965	21.91000	6.72541	5.01516	6.80017	1.71152	6.89451	0.72396
4	737.31802		7.95478	325.95893	8.00132	25.50477	8.07195	5.82011	8.16691	1.97742	8.28690	0.83127
5	861.64968		9.61839	380.39514	9.67856	29.70139	9.76943	6.75494	9.89175	2.28374	10.04654	0.95337
6	1008.5802		11.74323	444.54814	11.82178	34.62944	11.94052	7.84610	12.10056	2.63789	12.30342	1.09241
7	1183.6432		14.50032	520.73882	14.60466	40.45801	14.76254	9.12757	14.97563	3.04910	15.24624	1.25088
8	1394.3075		18.14238	612.07821	18.28368	47.41173	18.49776	10.64364	18.78717	3.52889	19.15548	1.43147
9	1650.8846		23.05215	722.82833	23.24788	55.79504	23.54473	12.45301	23.94684	4.09171	24.45984	1.63690
10	1967.9875		29.82774	858.97404	30.10584	66.03006	30.52844	14.63477	31.10210	4.75562	31.83608	1.86935

TABLE II: Polarisabilities of the H_2^+ molecular ion, in atomic units.

v	$L = 0$		$L = 1$		$L = 2$		$L = 3$		$L = 4$		$L = 5$	
	α_s		α_s	α_t	α_s	α_t	α_s	α_t	α_s	α_t	α_s	α_t
0	3.1687258		3.1783035	-0.8033729	3.1975081	-0.1931423	3.2264392	-0.0914467	3.2652493	-0.0544769	3.3141473	-0.0367142
1	3.8975634		3.9101018	-1.1442051	3.9352574	-0.2751013	3.9731892	-0.1302653	4.0241411	-0.0776138	4.0884471	-0.0523179
2	4.8215004		4.8380889	-1.6000689	4.8713900	-0.3847653	4.9216560	-0.1822373	4.9892726	-0.1086157	5.0747693	-0.0732474
3	6.0093275		6.0315483	-2.2129563	6.0761862	-0.5322759	6.1436389	-0.2521973	6.2345166	-0.1503892	6.3496578	-0.1014845
4	7.5604532		7.5906530	-3.0434869	7.6513642	-0.7322875	7.7432180	-0.3471422	7.8671844	-0.2071498	8.0246002	-0.1399110
5	9.6217735		9.6635170	-4.1811566	9.7475033	-1.0064626	9.8747452	-0.4774336	10.046804	-0.2851555	10.265837	-0.1928182
6	12.416000		12.474853	-5.7615823	12.593371	-1.3876723	12.773211	-0.6588274	13.016932	-0.3939491	13.328069	-0.2667729
7	16.290999		16.375936	-7.9965515	16.547168	-1.9273337	16.807463	-0.9160304	17.161118	-0.5485440	17.614095	-0.3721509
8	21.809473		21.935532	-11.228720	22.189990	-2.7087984	22.577626	-1.2892120	23.105870	-0.7734466	23.785138	-0.5259729
9	29.920328		30.113886	-16.036300	30.505195	-3.8730473	31.102847	-1.8465559	31.920266	-1.1104555	32.976407	-0.7574477
10	42.306330		42.616316	-23.445884	43.244200	-5.6711124	44.206257	-2.7100058	45.528094	-1.6347702	47.246181	-1.1195247

where the complex exponents α , β , γ , are generated in a pseudo-random way. The use of complex exponents instead of real ones allows reproducing the oscillatory behavior of the vibrational part of the wave function and improves the convergence rate. In numerical calculations we utilize basis sets as large as $N = 7\,000$ functions, in order to provide the required accuracy for the static polarisability of about 8 significant digits.

We note that a variational principle holds for the numerical value for α_s (but not for α_t): the larger the value, the closer it is to the exact (non-relativistic) value, provided that the initial wave function is accurate enough.

The results of numerical calculations of the polarisabilities for a wide range of ro-vibrational states are presented in Tables 1, 2, 3. These polarisabilities do not include relativistic corrections. These have so far been computed only for the ground rovibrational level ($v = 0$, $L = 0$) of H_2^+ [25]. Therefore, the relative inaccuracy of the values of the table as compared to the exact values is of order $\alpha^{-2} \simeq 1 \times 10^{-4}$. This is sufficiently small for current and near-future purposes.

C. Scaling with rotational angular momentum

For large L , we find for HD^+ ,

$$\alpha_t(v, L) \propto \frac{1}{L(L+1)(2L-1)(2L+3)}. \quad (12)$$

This follows from an argument described below after Eq. (25).

TABLE III: Polarisabilities of the D_2^+ molecular ion, in atomic units.

v	$L = 0$	$L = 1$		$L = 2$		$L = 3$		$L = 4$		$L = 5$	
	α_s	α_s	α_t	α_s	α_t	α_s	α_t	α_s	α_t	α_s	α_t
0	3.0719887	3.0765904	-0.7579521	3.0858052	-0.1813435	3.0996560	-0.0852443	3.1181777	-0.0503016	3.1414173	-0.0335048
1	3.5530258	3.5585822	-0.9782731	3.5697111	-0.2340592	3.5864444	-0.1100266	3.6088309	-0.0649271	3.6369364	-0.0432481
2	4.1195817	4.1263238	-1.2485988	4.1398301	-0.2987476	4.1601453	-0.1404432	4.1873367	-0.0828824	4.2214959	-0.0552137
3	4.7912827	4.7995087	-1.5808716	4.8159913	-0.3782716	4.8407920	-0.1778439	4.8740043	-0.1049671	4.9157545	-0.0699367
4	5.5933149	5.6034134	-1.9904009	5.6236531	-0.4763025	5.6541185	-0.2239603	5.6949390	-0.1322078	5.7462891	-0.0881048
5	6.5583187	6.5708021	-2.4970077	6.5958274	-0.5975951	6.6335113	-0.2810366	6.6840342	-0.1659357	6.7476365	-0.1106108
6	7.7290547	7.7446049	-3.1266348	7.7757864	-0.7483752	7.8227607	-0.3520126	7.8857801	-0.2078964	7.9651778	-0.1386263
7	9.1622096	9.1817469	-3.9136471	9.2209342	-0.9368936	9.2799934	-0.4407873	9.3592871	-0.2604068	9.4592730	-0.1737083
8	10.933925	10.958708	-4.9041723	11.008431	-1.1742306	11.083398	-0.5526002	11.184144	-0.3265836	11.311297	-0.2179539
9	13.147977	13.179752	-6.1610454	13.243527	-1.4754879	13.339760	-0.6946003	13.469145	-0.4106838	13.632624	-0.2742310
10	15.948121	15.989359	-7.7712809	16.072159	-1.8615919	16.197178	-0.8766990	16.365416	-0.5186172	16.578236	-0.3465280

v	$L = 0$	$L = 1$		$L = 2$		$L = 3$		$L = 4$	
	$\delta\alpha_s$	$\delta\alpha_s$	$\delta\alpha_t$	$\delta\alpha_s$	$\delta\alpha_t$	$\delta\alpha_s$	$\delta\alpha_t$	$\delta\alpha_s$	$\delta\alpha_t$
0	3.3	3.1	-0.72	3.1	-0.18	3.2	-0.08	3.2	-0.06
1	3.9	3.7	-0.98	3.8	-0.24	3.8	-0.12	3.8	-0.08
2	4.7	4.5	-1.34	4.5	-0.34	4.6	-0.16	4.6	-0.10
3	5.6	5.4	-1.80	5.5	-0.44	5.5	-0.22	5.6	-0.12
4	6.8	6.6	-2.42	6.6	-0.60	6.7	-0.28	6.8	-0.16

TABLE IV: Difference $\delta\alpha$ between the accurate polarisabilities of HD^+ (this work) and those computed by a summation over all intermediate rovibrational states in the ground electronic state, in atomic units. The latter are calculated from the results of Ref. [26] (which are there given in terms of $\alpha_{vLFSJJ_z}^{(t)}$, $\alpha_{vLFSJJ_z}^{(t)}$) as $\alpha_s(v, L) = (\alpha_{vLFSJJ_z}^{(t)} + 2\alpha_{vLFSJJ_z}^{(t)})/3$ (where any hyperfine state (F, S, J, J_z) can be chosen) and $\alpha_t(v, L) = (\alpha_{vLn_s}^{(t)} - \alpha_{vLn_s}^{(t)})/L(2L - 1)$, where n_s denotes the stretched state, defined in Sec. III C 3.

We have found heuristically, that for H_2^+ and D_2^+ ,

$$\alpha_t(v, L) \propto \frac{1}{(2L - 1)(2L + 3)}. \quad (13)$$

D. Comparison with previous work

1. Contribution from the ground electronic state

An approximation to the polarisability can be obtained using the well-known sum-over-intermediate-states expression, where the sum is truncated to a subset of levels. For HD^+ , such a calculation has been performed using transition dipole moments computed in the Born-Oppenheimer approximation [26], including in the sum only levels of the ground electronic state (the inaccuracy of the used dipole moments is mentioned further below). At first sight, it may appear that the polarisability of a level (v, L) in HD^+ is dominated by the contribution from the rotational levels adjacent in energy to the particular state, namely ($v, L \pm 1$). This is evidently true for $L = 0$ levels. However, for $L \neq 0$, there is partial cancellation of the two contributions from $L' = L \pm 1$. Even then, the α_t values indeed arise essentially from the rovibrational transitions. However, the $\alpha_s(L > 0)$ values are actually dominated by the contribution from the excited electronic states. The comparison of the accurate results given in the tables above with the truncated-sum results allows putting in evidence the contribution from the excited electronic states. The comparison is shown in Table IV, showing that for low-lying rovibrational levels ($v, L < 5$), the difference is of order several atomic units for α_s and less than 2.5 atomic units for α_t . The increase of the difference with v is due to the fact that the contributions from excited electronic states become more important since the level v is getting closer in energy to them.

For the homonuclear H_2^+ and D_2^+ the polarisability arises only from the excited electronic states, since there is no

electric-dipole coupling between levels of the ground electronic state. As a consequence, the polarisabilities $\alpha_s(L=0)$ and $\alpha_t(v, L)$ are much smaller than in the case of the heteronuclear ions, as has been noted in previous studies cited above.

2. General calculations

We can compare our results with some previous studies.

Early on, Bishop and Lam [19] studied the states $v=0, L=0-10$ of H_2^+ . The largest number of levels was considered by Moss and Valenzano, who covered the three ion species also treated here, with $L=0, 1$, and all v [21]. Our results agree with theirs, to within two units of the last digit reported by them, except for the level ($v=8, L=1$), where the largest discrepancy occurs, 0.007 at. u.

The agreement with the $L=0$ values for the three ion species determined by Hilico et al. [20], and Karr et al. [23] is better than 4×10^{-8} in relative terms.

Pilon and Baye recently computed the polarisabilities of H_2^+ for a number of levels [27]. The values for $v=0, L=0, 1, 2, 3, 4, 5$ agree to better than 2×10^{-8} in relative terms. For $L=1, v=0, 1, 2, 3$ and for $L=2, v=0, 1, 2, 3$ the values agree with the present values to better than 3×10^{-7} in relative terms.

III. PERTURBATION THEORY FOR THE HYPERFINE STATES

A. Energy shifts

The hyperfine interactions split each rovibrational level into a number of hyperfine sub-levels. We denote the corresponding kets as $|m\rangle = |v L n J_z\rangle$, where n is a label for the particular hyperfine state in a rovibrational level (v, L) (note that this notation includes both pure and non-pure spin states). n is written as (F, S, J) for HD^+ and (I, S, J) for H_2^+ , see Sec. IIIB below. When the Stark shifts of the quantum levels are small compared to other shifts, we can apply first-order perturbation theory. The Stark energy shift of a state $|m\rangle$ can be expressed in different ways (for simplicity, in the following we omit the caret on the polarization operators) [28]:

$$\begin{aligned} \Delta E(m) &= -\frac{1}{2} [\langle m|\alpha_{\parallel}|m\rangle E_z^2 + \langle m|\alpha_{\perp}|m\rangle (E_x^2 + E_y^2)] , \\ &= -\frac{1}{2} \left[\alpha_s(E_x^2 + E_y^2 + E_z^2) + \alpha_t(E_z^2 - \frac{1}{2}(E_x^2 + E_y^2)) \langle m|\sqrt{\frac{8}{3}}\{\mathbf{L} \otimes \mathbf{L}\}_{20}|m\rangle \right] , \\ &= -\frac{1}{2} \alpha_s(E_x^2 + E_y^2 + E_z^2) - \alpha_t(E_z^2 - \frac{1}{2}(E_x^2 + E_y^2)) \langle m|L_z^2 - \frac{1}{3}\mathbf{L}^2|m\rangle , \\ &= -\frac{1}{2} \mathbf{E}^2 \left(\alpha_s + \alpha_t(3 \cos^2 \theta - 1) \langle m|L_z^2 - \frac{1}{3}\mathbf{L}^2|m\rangle \right) , \\ &= -\frac{1}{2} \mathbf{E}^2 \left(\alpha_s(v, L) + (3 \cos^2 \theta - 1) \frac{1}{3} \langle v L n J_z|\alpha_{\parallel} - \alpha_{\perp}|v L n J_z\rangle \right) , \end{aligned} \quad (14)$$

where θ is the angle between the quantization axis and the direction of the electric field $\mathbf{E}$.

In Refs. [16, 26] the levels shifts were described in terms of longitudinal polarisability, $\alpha^{(l)}$, and transverse polarisability $\alpha^{(t)}$. They are related to the expectation values of the operators introduced here by $\langle m|\alpha_{\parallel}|m\rangle = \alpha^{(l)}$ and $\langle m|\alpha_{\perp}|m\rangle = \alpha^{(t)}$.

B. Hyperfine structure

We limit ourselves in the following to the ion species H_2^+ and HD^+ , which are most relevant for experimental work at present.

In case of the molecular ion H_2^+ we have identical nuclei and nuclear permutation symmetry. This makes some spin configurations forbidden and splits the consideration of hyperfine states into two cases (see [29]): for even L , the total nuclear spin I is zero and only two hyperfine sub-levels are possible; for states with odd L , the total nuclear spin is one and the ro-vibrational level is split into 5 or 6 hyperfine sub-levels, depending on the value of L .

The most suitable coupling scheme of angular momentum operators is

$$\mathbf{S} = \mathbf{I} + \mathbf{s}_e, \quad \mathbf{J} = \mathbf{S} + \mathbf{L}, \quad (15)$$

where $\mathbf{I}$ is the total nuclear spin operator, and $\mathbf{s}_e$ is the electron spin operator. The basis states which correspond to this coupling are

$$|ISLJJ_z\rangle = \sum_{I_z, \xi, S_z} C_{SS_z, L L_z}^{JJ_z} C_{I I_z, s_e \xi}^{SS_z} \left(|I I_z\rangle \cdot |s_e \xi\rangle \cdot |L L_z\rangle \right), \quad (16)$$

and will be called pure states [30].

The effective HFS Hamiltonian is expressed as [29]

$$\begin{aligned} H_{\text{eff}} = & b_F(\mathbf{I} \cdot \mathbf{s}_e) + c_e(\mathbf{L} \cdot \mathbf{s}_e) + c_I(\mathbf{L} \cdot \mathbf{I}) \\ & + \frac{d_1}{(2L-1)(2L+3)} \left\{ \frac{2}{3} \mathbf{L}^2(\mathbf{I} \cdot \mathbf{s}_e) - [(\mathbf{L} \cdot \mathbf{I})(\mathbf{L} \cdot \mathbf{s}_e) + (\mathbf{L} \cdot \mathbf{s}_e)(\mathbf{L} \cdot \mathbf{I})] \right\} \\ & + \frac{d_2}{(2L-1)(2L+3)} \left[\frac{1}{3} \mathbf{L}^2 \mathbf{I}^2 - \frac{1}{2}(\mathbf{L} \cdot \mathbf{I}) - (\mathbf{L} \cdot \mathbf{I})^2 \right]. \end{aligned} \quad (17)$$

For the case of even L , the pure states are the true HFS eigenstates, since the 2×2 effective HFS Hamiltonian matrix is diagonal. Even for odd- L states, the pure states are good approximations to the true HFS states [30], since the coefficients of admixture of other states to a given true HFS state are small, e.g. for $L = 1$ do not exceed 0.04, and for $L = 3$ do not exceed 0.06. This means that even in this case, a good approximation for expectation values such as Eq. (14) may be obtained using the pure states.

For the hydrogen molecular ion HD^+ the coupling scheme of the particle angular momentum operators is [31]

$$\mathbf{F} = \mathbf{I}_p + \mathbf{s}_e, \quad \mathbf{S} = \mathbf{F} + \mathbf{I}_d, \quad \mathbf{J} = \mathbf{S} + \mathbf{L}. \quad (18)$$

$\mathbf{I}_p$, $\mathbf{I}_d$ are the proton and deuteron spin operators, respectively. The effective Hamiltonian is given in Ref. [31]. The pure states are determined in a similar way as in Eq. (16). In zero magnetic field, the pure states represent a good approximation to some of the true HFS states, and may be used to calculate approximate values of the polarisabilities. Details are given in Sec. IV below. Hyperfine states are labeled by $n = (F S J)$.

C. Analytical results

In this subsection we discuss some useful results that allow to understand several dependencies. In particular we discuss the polarisabilities of the pure spin states, for two reasons. First, a significant part of hyperfine states may be well approximated by pure spin states; second, since all hyperfine states can be expressed as weighted sums of pure spin states, their polarisabilities can conveniently be computed from the pure state polarisabilities.

1. Zero magnetic field

When the magnetic field is zero, the total angular momentum squared $\mathbf{J}^2$ commutes with the hyperfine Hamiltonian and J is a good quantum number. Therefore we can apply the Wigner-Eckart theorem, and separate the J_z -dependence of the expectation value:

$$\begin{aligned} \langle vLnJ_z | \{\mathbf{L} \otimes \mathbf{L}\}_{20} | vLnJ_z \rangle &= C_{20, JJ_z}^{JJ_z} \langle vLn | \{\mathbf{L} \otimes \mathbf{L}\}_2 | vLn \rangle / \sqrt{2J+1} \\ &= - \frac{J(J+1) - 3J_z^2}{\sqrt{J(J+1)(2J-1)(2J+1)(2J+3)}} \langle vLn | \{\mathbf{L} \otimes \mathbf{L}\}_2 | vLn \rangle \end{aligned} \quad (19)$$

We therefore obtain the J_z - dependence of the polarisability anisotropy as follows:

$$\langle vLnJ_z|\alpha_{\parallel} - \alpha_{\perp}|vLnJ_z\rangle = \left(1 - \frac{3J_z^2}{J(J+1)}\right) \langle vLnJ_z=0|\alpha_{\parallel} - \alpha_{\perp}|vLnJ_z=0\rangle. \quad (20)$$

Note that this result holds both for pure and non-pure spin states. It follows that for $J = 0$ states, the polarisability anisotropy is zero. For HD^+ , $J = 0$ states can only occur for $L < 3$, since the minimum J value permitted by angular momentum algebra is $L - 2$. For H_2^+ , there are no such states, since J is a half-integer number.

2. Pure states

For pure angular momentum states, the matrix elements of the polarisability anisotropy can be evaluated explicitly. Considering only the coupling scheme $\mathbf{J} = \mathbf{S} + \mathbf{L}$, we have (note that this is independent of I or F)

$$\begin{aligned} \left\langle SLJJ_z \left| 2 \left(L_z^2 - \frac{1}{3} \mathbf{L}^2 \right) \right| SLJJ_z \right\rangle &= \sqrt{\frac{8}{3}} \langle SLJJ_z | \{\mathbf{L} \otimes \mathbf{L}\}_{20} | SLJJ_z \rangle, \\ &= \sqrt{\frac{8}{3}} \frac{C_{JJ_z, 20}^{JJ_z}}{\sqrt{2J+1}} \langle SLJ \| \{\mathbf{L} \otimes \mathbf{L}\}_2 \| SLJ \rangle, \\ &= -\frac{(J(J+1) - 3J_z^2) [3D(D-1) - 4J(J+1)L(L+1)]}{3J(J+1)(2J-1)(2J+3)}, \end{aligned} \quad (21)$$

where

$$D = J(J+1) + L(L+1) - S(S+1). \quad (22)$$

This result is obtained using the following relations [32, 33]

$$\begin{aligned} \langle SLJ \| \{\mathbf{L} \otimes \mathbf{L}\}_2 \| SLJ \rangle &= (2J+1)(-1)^{S+L+J+2} \left\{ \begin{matrix} L & L & 2 \\ J & J & S \end{matrix} \right\} \langle L \| \{\mathbf{L} \otimes \mathbf{L}\}_2 \| L \rangle, \\ \langle L \| \{\mathbf{L} \otimes \mathbf{L}\}_2 \| L \rangle &= \frac{1}{\sqrt{6}} \sqrt{L(L+1)} \sqrt{(2L-1)(2L+1)(2L+3)}. \end{aligned}$$

In H_2^+ we consider first the states having even L , so $I = 0$. Then $S = 1/2$. These pure states are exact HFS eigenstates, and therefore Eq. (21) immediately gives the exact Stark shift using Eqs. (14, 20):

$$\langle m(\text{even } L) | \alpha_{\parallel} - \alpha_{\perp} | m(\text{even } L) \rangle_{\text{H}_2^+} = -\frac{3}{2} \alpha_t(v, L) \frac{(J(J+1) - 3J_z^2) [3D(D-1) - 4J(J+1)L(L+1)]}{3J(J+1)(2J-1)(2J+3)}. \quad (23)$$

with $D = J(J+1) + L(L+1) - 3/4$.

For pure states with odd L (and therefore $I = 1$):

$$\begin{aligned} \left\langle I = 1 SLJ \left\| \{\mathbf{L} \otimes \mathbf{L}\}_2 \right\| I = 1 SLJ \right\rangle &= (2J+1)(-1)^{S+L+J} \left\{ \begin{matrix} L & L & 2 \\ J & J & S \end{matrix} \right\} \langle L \| \{\mathbf{L} \otimes \mathbf{L}\}_2 \| L \rangle \\ &= \sqrt{\frac{2J+1}{6}} \frac{3D(D-1) - 4J(J+1)L(L+1)}{2\sqrt{J(J+1)(2J-1)(2J+3)}}, \end{aligned} \quad (24)$$

where D is given by Eq. (22). We see that the actual value of I does not occur on the r.h.s., and that we obtain the same result as for the $I = 0$ pure states. Eq. (23) is an approximate result also for the odd- L hyperfine states of H_2^+ which are not pure, provided they are approximately pure (see below).

In the case of HD^+ , where the pure states are denoted as $|FSLJ J_z\rangle$, Eq. (23) also holds, where L now can be even or odd. There is no dependence on F .

Summarizing, for any pure state of H_2^+ and HD^+ , and, by consequence, also for all other molecular hydrogen ions, Eq. (23) gives the polarisability anisotropy:

$$\langle \text{pure state} | \alpha_{\parallel} - \alpha_{\perp} | \text{pure state} \rangle_{\text{any species}} = -\frac{3}{2} \alpha_t(v, L) \frac{(J(J+1) - 3J_z^2) [3D(D-1) - 4J(J+1)L(L+1)]}{3J(J+1)(2J-1)(2J+3)}. \quad (25)$$

We note that Roeggen [34] has developed an approximate theory of the polarisability of heteronuclear diatomic molecules with spin, neglecting nuclear spin. If we combine our Eq. (25) and the approximate dependencies Eq. (12) we reproduce the result given in Eqs. (62, 63) of Ref. [34].

3. The stretched states

The stretched states are those exact HFS states having maximal total angular momentum J and maximal (absolute) projection $|J_z|$. These are also pure states. In HD^+ , these are the states $|v L n_s\rangle$, where n_s denotes the stretched hyperfine state: $F = 1$, $S = 2$, $J = L + 2$, $J_z = \pm(L + 2)$. We find from Eq. (23) or by analytical evaluation of the matrix elements for these two stretched states (the evaluation is simple, if the calculations is done with the basis functions being the eigenfunctions of the individual angular momenta I_p, I_d, S, L),

$$\langle v L n_s | \alpha_{\parallel} - \alpha_{\perp} | v L n_s \rangle = L(2L - 1) \alpha_t(v, L). \quad (26)$$

Compare the discussion in Ref. [16].

In H_2^+ , the stretched states are $|v L n_s\rangle \equiv |v, L, I = 1, S = 3/2, J = L + 3/2, J_z = \pm(L + 3/2)\rangle$. The same result Eq. (26) is obtained.

By evaluating the polarisabilities of all hyperfine states, we find that if $L > 1$, the largest value of $\langle m | \alpha_{\parallel} - \alpha_{\perp} | m \rangle$ within a rovibrational level occurs for the stretched states (see tables below). Therefore, in the following discussion, we normalize the polarisability anisotropy values of any hyperfine state in a particular rovibrational level relative to that of the stretched states in that same level.

For HD^+ , combining the result Eq. (26) for the stretched states with the approximate behavior Eq. (12), we obtain $\langle v L n_s | \alpha_{\parallel} - \alpha_{\perp} | v L n_s \rangle \propto (3 + 5L + 2L^2)^{-1}$. This describes a rather strong decrease in the magnitude of all anisotropic polarisability values, not only those of the stretched-states, with increasing L .

IV. NUMERICAL RESULTS FOR THE HYPERFINE-STATE DEPENDENCE

The evaluation of the matrix elements of Eq. (10) for all (exact) hyperfine states is straightforward, once the hyperfine states in absence of electric field are known. The calculation can for example proceed by considering the expansion of the hyperfine states in pure states, and then applying Eq. (25), which holds for the pure states of any molecular hydrogen ion. Actually, the matrix elements are the same (apart from prefactors such as α_t) as the matrix elements for the electric quadrupole shift evaluated in Ref. [16], and an explicit formula is given there.

We have performed the computation for the rovibrational levels up to $v = 4$ and $L = 4$. Note that the polarisability anisotropy vanishes for $L = 0$ states and is therefore not reported in the tables. We confine ourselves to the case of zero magnetic field.

The results are summarized in table V and table VI where we give the values for the hyperfine states having $J_z = 0$. The values for $J_z \neq 0$ can be easily obtained using Eq. (20). Note that for a given hyperfine state and value of L the dependence on v is usually weak, limited to several percent, except for a few cases.

By looking at the values in the Tables V, VI, one can see that the approximation that the polarisability does not depend on F is quite good for some hyperfine states, and moderate in others, which is due to their more or less pure character. In order to obtain values accurate to better than one atomic unit for HD^+ , because of its large values of α_t it is necessary to use the exact hyperfine dependence of the polarisability anisotropy,

The results for H_2^+ in odd - L states are shown in Table VII. We can see that in this species, the anisotropic polarisabilities are always very close to those of the pure states. The maximum deviation is approximately 0.01 atomic unit. Thus, for current purposes, for H_2^+ one may use Eq. (25) for all rovibrational levels.

hyp. state (F, S, J, J_z)	Eq. (25) (normalized)	$(v, L) = (0, 1)$	$(v, L) = (1, 1)$	level $(v, L) = (2, 1)$	$(v, L) = (3, 1)$	$(v, L) = (4, 1)$
(0, 1, 2, 0)	-1	-0.999422	-0.999457	-0.999491	-0.999525	-0.999558
(0, 1, 1, 0)	1	0.998747	0.998825	0.998902	0.998978	0.999053
(0, 1, 0, 0)*	0	0	0	0	0	0
(1, 0, 1, 0)	-2	-1.66845	-1.69239	-1.71585	-1.73892	-1.76151
(1, 1, 1, 0)	1	0.590503	0.619326	0.647628	0.675521	0.702883
(1, 1, 0, 0)*	0	0	0	0	0	0
(1, 1, 2, 0)	-1	-0.999973	-0.999972	-0.999971	-0.99997	-0.99997
(1, 2, 1, 0)	-0.2	-0.120803	-0.125763	-0.130681	-0.135574	-0.140428
(1, 2, 3, 0)*	-0.8	-0.8	-0.8	-0.8	-0.8	-0.8
(1, 2, 2, 0)	1	0.999395	0.999429	0.999462	0.999495	0.999528

TABLE V: The normalized anisotropic polarisabilities for the $J_z = 0$ - hyperfine states of HD^+ in $L = 1$ levels. The first column shows the hyperfine state's label n, J_z (F and S are usually approximate, J is an exact quantum number), the second column contains the normalized values $\langle v L F S J J_z = 0 | \alpha_{\parallel} - \alpha_{\perp} | v L F S J J_z = 0 \rangle / \langle v L n_s | \alpha_{\parallel} - \alpha_{\perp} | v L n_s \rangle$ for the pure state $|v L F S J J_z = 0\rangle$ giving the largest contribution to the exact HFS state $|v L n J_z = 0\rangle$ (note that the values are independent of F and of v). The following columns give the actual values $\langle v L n J_z = 0 | \alpha_{\parallel} - \alpha_{\perp} | v L n J_z = 0 \rangle / \langle v L n_s | \alpha_{\parallel} - \alpha_{\perp} | v L n_s \rangle$ for each hyperfine state. The normalization is with respect to the polarisability anisotropy of the stretched state $|v L n_s\rangle$ of the same rovibrational level. The three states marked with an asterisk are pure states.

hyp. state (F, S, J, J_z)	Eq. (25) (normalized)	$(v, L) = (0, 2)$	$(v, L) = (1, 2)$	level(v, L) $(v, L) = (2, 2)$	$(v, L) = (3, 2)$	$(v, L) = (4, 2)$
(0, 1, 3, 0)	-0.8	-0.799487	-0.799517	-0.799548	-0.799578	-0.799608
(0, 1, 2, 0)	-0.5	-0.499349	-0.49939	-0.49943	-0.499469	-0.499508
(0, 1, 1, 0)	-0.7	-0.698944	-0.699011	-0.699076	-0.699141	-0.699205
(1, 0, 2, 0)	-1	-0.89668	-0.902108	-0.907625	-0.913227	-0.918913
(1, 1, 1, 0)	-0.7	-0.598522	-0.605382	-0.612152	-0.618825	-0.625393
(1, 1, 2, 0)	-0.5	-0.572569	-0.56875	-0.564853	-0.560883	-0.556839
(1, 1, 3, 0)	-0.8	-0.798847	-0.798904	-0.798961	-0.799019	-0.799079
(1, 2, 0, 0)*	0	0	0	0	0	0
(1, 2, 1, 0)	0.7	0.597466	0.604392	0.611229	0.617966	0.624598
(1, 2, 2, 0)	0.2143	0.182884	0.184534	0.186194	0.187865	0.189546
(1, 2, 3, 0)	-0.2	-0.201666	-0.201579	-0.201491	-0.201403	-0.201313
(1, 2, 4, 0)*	-0.7143	-0.714286	-0.714286	-0.714286	-0.714286	-0.714286
		$(v, L)=(0,3)$	$(v, L)=(1,3)$	$(v, L)=(2,3)$	$(v, L)=(3,3)$	$(v, L)=(4,3)$
(0, 1, 4, 0)	-0.7143	-0.713807	-0.713835	-0.713864	-0.713892	-0.71392
(0, 1, 3, 0)	-0.6	-0.599383	-0.599421	-0.599459	-0.599496	-0.599533
(0, 1, 2, 0)	-0.6857	-0.684872	-0.684925	-0.684977	-0.685028	-0.685079
(1, 0, 3, 0)	-0.8	-0.743487	-0.745805	-0.748206	-0.750693	-0.753272
(1, 1, 2, 0)	-0.6857	-0.641563	-0.644157	-0.646756	-0.649358	-0.651959
(1, 1, 3, 0)	-0.6	-0.638925	-0.637509	-0.636019	-0.634449	-0.632793
(1, 1, 4, 0)	-0.7143	-0.712034	-0.712155	-0.712277	-0.712399	-0.712523
(1, 2, 1, 0)*	-0.48	-0.48	-0.48	-0.48	-0.48	-0.48
(1, 2, 2, 0)	-0.1714	-0.216421	-0.213776	-0.211124	-0.208471	-0.205819
(1, 2, 3, 0)	-0.2533	-0.271538	-0.270598	-0.26965	-0.268696	-0.267735
(1, 2, 4, 0)	-0.4286	-0.431302	-0.431152	-0.431002	-0.430852	-0.4307
(1, 2, 5, 0)*	-0.6667	-0.666667	-0.666667	-0.666667	-0.666667	-0.666667
		$(v, L)=(0,4)$	$(v, L)=(1,4)$	$(v, L)=(2,4)$	$(v, L)=(3,4)$	$(v, L)=(4,4)$
(0, 1, 5, 0)	-0.6667	-0.666213	-0.66624	-0.666267	-0.666293	-0.66632
(0, 1, 4, 0)	-0.6071	-0.606572	-0.606607	-0.606642	-0.606676	-0.60671
(0, 1, 3, 0)	-0.6548	-0.654035	-0.654081	-0.654126	-0.65417	-0.654214
(1, 0, 4, 0)	-0.7143	-0.677583	-0.678822	-0.68012	-0.68148	-0.682906
(1, 1, 3, 0)	-0.6548	-0.627904	-0.62934	-0.630793	-0.632265	-0.633749
(1, 1, 4, 0)	-0.6071	-0.630532	-0.629989	-0.62939	-0.628733	-0.628016
(1, 2, 2, 0)*	-0.5612	-0.561224	-0.561224	-0.561224	-0.561224	-0.561224
(1, 1, 5, 0)	-0.6667	-0.663695	-0.663856	-0.664019	-0.664183	-0.664348
(1, 2, 3, 0)	-0.3929	-0.420442	-0.41896	-0.417462	-0.415946	-0.414418
(1, 2, 4, 0)	-0.4096	-0.423439	-0.422708	-0.421974	-0.421236	-0.420494
(1, 2, 5, 0)	-0.5	-0.503426	-0.503237	-0.503047	-0.502857	-0.502666
(1, 2, 6, 0)*	-0.6364	-0.636364	-0.636364	-0.636364	-0.636364	-0.636364

TABLE VI: Same as Table V, but for levels $L = 2, 3, 4$. Note that the values in the second column are rounded.

hyp. state (I, S, J, J_z)	Eq. (25) (normalized)	$v = 0$	$v = 1$	level (v, L) $v = 2$	$v = 3$	$v = 4$
				$L = 1$		
$(1, \frac{1}{2}, \frac{3}{2}, 0)$	-1.25	-1.24946	-1.2495	-1.24951	-1.24956	-1.2496
$(1, \frac{1}{2}, \frac{1}{2}, 0)$	-0.125	-0.1216	-0.12186	-0.12212	-0.12238	-0.12262
$(1, \frac{3}{2}, \frac{1}{2}, 0)$	2.125	2.12158	2.12185	2.12212	2.12239	2.12262
$(1, \frac{3}{2}, \frac{5}{2}, 0)^*$	-0.875	-0.875	-0.875	-0.875	-0.875	-0.875
$(1, \frac{3}{2}, \frac{3}{2}, 0)$	1.	0.99946	0.9995	0.99952	0.99956	0.9996
				$L = 3$		
$(1, \frac{1}{2}, \frac{7}{2}, 0)$	-0.75	-0.74955	-0.74958	-0.74961	-0.74964	-0.74967
$(1, \frac{1}{2}, \frac{5}{2}, 0)$	-0.75	-0.74871	-0.7488	-0.7489	-0.74899	-0.74909
$(1, \frac{3}{2}, \frac{3}{2}, 0)^*$	-0.6	-0.6	-0.6	-0.6	-0.6	-0.6
$(1, \frac{3}{2}, \frac{5}{2}, 0)$	-0.4125	-0.41379	-0.41369	-0.41359	-0.4135	-0.41341
$(1, \frac{3}{2}, \frac{7}{2}, 0)$	-0.5	-0.50044	-0.50041	-0.50038	-0.50035	-0.50032
$(1, \frac{3}{2}, \frac{9}{2}, 0)^*$	-0.6875	-0.6875	-0.6875	-0.6875	-0.6875	-0.6875

TABLE VII: Anisotropic polarisability of the $J_z = 0$ - quantum states of H_2^+ in $L = 1, 3$, normalized to those of the stretched states. See caption of Tab. V.

v	L	$\Delta E_{stat}((v, L), T)$ [mHz]	$\Delta E_{dyn,elec}((v, L), T)$ [mHz]
0	0	-27.3	-0.0023
0	1	-27.4	-0.0023
0	3	-27.8	-0.0024
1	1	-33.7	-0.0044
1	3	-34.2	-0.0046
2	1	-41.7	
3	1	-51.9	-0.0156

TABLE VIII: Static approximation of the BBR shift and dynamic contribution of some levels of H_2^+ , at $T = 300$ K. The total BBR shift is obtained by adding the values in the third and fourth columns.

V. THE BLACK-BODY RADIATION FREQUENCY SHIFT

A. Generalities

The black-body radiation (BBR) shift of a level m is computed as

$$\Delta E_{BBR}(m, T) = -\frac{1}{2} \int_0^\infty \alpha_s(m, \omega) \mathcal{E}_{BBR}(T, \omega)^2 d\omega. \quad (27)$$

if the BBR electric field is unpolarized. The contributions from the magnetic field are neglected. Therefore, under this assumption and because of the small hyperfine splittings compared to the (smallest) rotational levels splitting (20 MHz versus 1 THz, i.e. 2×10^{-5} in relative terms), the BBR shift is to a high approximation equal for all hyperfine states of a given rovibrational level.

B. Approximate treatment

1. Homonuclear ions

We may approximate the polarisability of the homonuclear ions by its zero-frequency value: $\alpha_s((v, L), \omega) \simeq \alpha_s((v, L), \omega = 0) = \alpha_s(v, L)$, where the values are given in the Tables II and III above. Then

$$\Delta E_{BBR}(m, T) \simeq \Delta E_{stat}((v, L), T) = -\frac{1}{2} \alpha_s((v, L), 0) (831.9 \text{ V/m})^2 (T/300 \text{ K})^4. \quad (28)$$

(In this expression, the value of α_s in atomic units is to be multiplied by the value of $4\pi\epsilon_0 a_0^3$ in SI units). A polarisability of 1 atomic unit gives a frequency shift of -8.6 mHz at 300 K. The shifts of several selected rovibrational levels are given in Table VIII.

2. Heteronuclear ions

For the heteronuclear ions, we express the polarisability as

$$\alpha_s((v, L), \omega) = \alpha_s((v, L), 0) + \delta\alpha_{s,dyn,rv}((v, L), \omega) + \delta\alpha_{s,dyn,elec}((v, L), \omega). \quad (29)$$

Here, $\delta\alpha_{s,dyn,elec}((v, L), \omega)$ is the frequency-dependent contribution from the excited electronic levels. It does not include the frequency-independent part, which is instead included in $\alpha_s((v, L), 0)$. Both $\delta\alpha_{s,dyn,elec}((v, L), \omega)$ and $\delta\alpha_{s,dyn,rv}((v, L), \omega)$ are defined so that they vanish at $\omega = 0$. The frequency-dependent contributions from E1

TABLE IX: Selected reduced transition dipole matrix elements, $d = \langle v' L' | \mathbf{d} | v L \rangle$, for transitions between rovibrational states of HD^+ for the case $L = 5 \rightarrow L' = 6$ (in atomic units). The notation $[x]$ means $\times 10^x$.

$v \rightarrow v'$	d	$v \rightarrow v'$	d
$(0 \rightarrow 0)$	0.85382285	$(3 \rightarrow 0)$	0.22598012[-02]
$(0 \rightarrow 1)$	0.64170498[-01]	$(3 \rightarrow 1)$	0.19539417[-01]
$(0 \rightarrow 2)$	0.97751608[-02]	$(3 \rightarrow 2)$	0.20027036
$(0 \rightarrow 3)$	0.23604947[-02]	$(3 \rightarrow 3)$	1.00176877
$(0 \rightarrow 4)$	0.74298137[-03]	$(3 \rightarrow 4)$	0.12862769
$(0 \rightarrow 5)$	0.27963915[-03]	$(3 \rightarrow 5)$	0.30336340[-01]
$(1 \rightarrow 0)$	0.11178434	$(4 \rightarrow 0)$	0.63859280[-03]
$(1 \rightarrow 1)$	0.90139089	$(4 \rightarrow 1)$	0.45000014[-02]
$(1 \rightarrow 2)$	0.90803829[-01]	$(4 \rightarrow 2)$	0.27891731[-01]
$(1 \rightarrow 3)$	0.16798347[-01]	$(4 \rightarrow 3)$	0.23541895
$(1 \rightarrow 4)$	0.46309897[-02]	$(4 \rightarrow 4)$	1.05507688
$(1 \rightarrow 5)$	0.16105915[-02]	$(4 \rightarrow 5)$	0.14394208
$(2 \rightarrow 0)$	0.11198057[-01]	$(5 \rightarrow 0)$	0.22302649[-03]
$(2 \rightarrow 1)$	0.16073279	$(5 \rightarrow 1)$	0.14049130[-02]
$(2 \rightarrow 2)$	0.95063090	$(5 \rightarrow 2)$	0.71058396[-02]
$(2 \rightarrow 3)$	0.11129675	$(5 \rightarrow 3)$	0.36420347[-01]
$(2 \rightarrow 4)$	0.23608027[-01]	$(5 \rightarrow 4)$	0.26815005
$(2 \rightarrow 5)$	0.72022091[-02]	$(5 \rightarrow 5)$	1.11088687

rovibrational transitions within the ground electronic state are important and give rise to [18],

$$\delta\alpha_{s,dyn,rv}((v, L), \omega) = \frac{1}{3} \frac{1}{2L+1} \sum_{v', L'} |\langle v' L' | \mathbf{d} | v, L \rangle|^2 \left(\frac{1}{E_{v' L'} - E_{v L} + \hbar\omega} + \frac{1}{E_{v' L'} - E_{v L} - \hbar\omega} - \frac{2}{E_{v' L'} - E_{v L}} \right). \quad (30)$$

In this sum, the value of L' can only take on the values $L \pm 1$, due to the selection rule. Again, $\alpha_s((v, L), 0)$ is the variational calculation result. As a first approximation, we can neglect $\delta\alpha_{s,dyn,elec}$, as done above for the homonuclear ions, since the transitions to the excited electronic states are of similar character. This neglect will be corrected in the next subsection.

The total BBR shift is

$$\Delta E_{BBR}((v, L), T) = \Delta E_{stat}((v, L), T) + \Delta E_{dyn,rv}((v, L), T) + \Delta E_{dyn,elec}((v, L), T). \quad (31)$$

We first discuss the dynamic *rovibrational* contribution to the BBR shift,

$$\Delta E_{dyn,rv}((v, L), T) = -\frac{1}{2} \int_0^\infty \delta\alpha_{s,dyn,rv}((v, L), \omega) \mathcal{E}_{BBR}(T, \omega)^2 d\omega,$$

which we have computed for levels up to $v_{max} = 10$, $L_{max} = 5$, extending the results of Ref. [18], which considered levels with $v_{max} = 7$, $L_{max} = 1$.

In this computation, it is important to use the most accurate transition dipoles values available, in order to reach a sufficient absolute accuracy in the polarisability and BBR shift, since partial cancellations occur in Eq. (30). For $v < 6$, $L < 6$ we use the precise transition dipoles of Tian et al. [35], based on variational wave functions. Their fractional inaccuracy is stated as smaller than 1×10^{-6} , and is less than that of our previously published values in Ref. [14]. As a check, we have recomputed the transition dipole moment of $(v = 0, L = 0) \rightarrow (v' = 0, L' = 1)$ with a larger basis set, and the value 0.3428334 at. u. in agreement with Tian et al. to better than 3×10^{-7} in fractional terms. We have computed the transition dipole moments between $L = 5$ and $L' = 6$ levels having $v, v' < 6$ in order to extend the results of Tian et al. They are listed in Tab. IX. For larger $v, v' \geq 6$ we use the Born-Oppenheimer transition dipole elements given in Ref. [26]. These agree, in the v, L range computed by Tian et al., within 1 to




FIG. 1: . The scalar polarisability of HD^+ in the level $(v = 1, L = 1)$, computed using variational wave functions. Atomic units are used.

2 parts in 10^4 with their results. As energy differences $E_{v,L} - E_{v',L'}$ we use the precise energies including QED corrections [36, 37] when $v < 5$, $L < 5$, and otherwise the values of Moss [38].

Table X (a) presents the relative value of the dynamic rovibrational contribution. We see that for the $L = 0$ levels a strong cancellation between the (particularly large) contributions $\Delta E_{stat}((v, L = 0), T)$ and $\Delta E_{dyn,rv}((v, L = 0), T)$ occurs, which results in a small BBR shift. In absolute terms, the BBR shift value is seen to grow with v and with L ; see part (b) of the Table. The absolute values are in the range of 1 mHz to several tens of mHz, for $v = 0 \dots 6$ and moderate L .

For $v < 6$ we estimate the inaccuracy of $\Delta E_{dyn,rv}((v, L), 300 \text{ K})$ to be less than 10^{-7} Hz, since the individual contributions to the sum are less than 0.1 Hz in absolute value. The values of $\Delta E_{stat}((v, L), 300 \text{ K})$ are smaller than 0.1 Hz in absolute value, and their inaccuracy is determined by the inaccuracy of our $\alpha_s((v, L), \omega = 0)$ values. The inaccuracy is thus less than 10^{-6} Hz. However, the non-relativistic approximation implies that both $\alpha_s((v, L), \omega = 0)$ and the transition dipoles are only accurate to the 1×10^{-4} fractional level. Then, the theoretical inaccuracy of the BBR shift, assuming the last term in Eq. (31) is negligible, may be stated conservatively as less than 3×10^{-5} Hz for the levels $v < 6$, since the shift and its uncertainty is mostly determined by three contributions, each with approximate uncertainty of 1×10^{-5} Hz. For $v \geq 6$, taking into account that the transition dipoles values are calculated in Born-Oppenheimer approximation, the overall inaccuracy is estimated at 6×10^{-5} Hz.

From an experimental point of view, the temperature derivative of the BBR shift is an important quantity, since the temperature of the BBR field in an ion trap has a relatively large uncertainty, due to the difficulty in determining it experimentally. For H_2^+ this derivative can be trivially obtained from Eq. (28), while the results for HD^+ are given in Tab. X (c, d). We find a strong variation between levels. Only for levels having larger v and L the normalized derivative is close to the value $4/T$ corresponding to a purely static BBR shift, Eq. (28).

C. Variational results

For several levels of both H_2^+ and HD^+ we have computed the dynamic polarisability $\alpha_{s,var}(\omega)$ (and $\alpha_{t,var}(\omega)$) *directly*, using variational wave functions. For one particular level of HD^+ , the polarisability α_s has been computed up to large frequencies, see Fig. 1. The calculation was performed using the complex coordinate rotation method [39, 40]. This overview clearly shows the dominating contributions from the rovibrational levels when ω is small whereas for large ω the excited electronic states yield a broad dispersive resonances. The low-frequency tail of this resonance, as $\omega \rightarrow 0$, is responsible for giving rise to $\Delta E_{dyn,elec}$.

For several other levels, the computation was performed up to an angular frequency $\omega = 0.1$ atomic units, in steps of 10^{-5} atomic units. The results are given in the additional material available online [41]. Since the computation was done in the non-relativistic approximation, the fractional inaccuracy of the values with respect to the exact values is approximately 1×10^{-4} . This is then also the fractional inaccuracy of the BBR shifts computed from this data.

1. H_2^+

For H_2^+ , we can compare our values of the scalar polarisability with the calculation by Pilon, who has communicated the values at six different frequencies [42]. The values agree, with deviations of at most 2×10^{-6} atomic units in the

(a)	v	L					
		0	1	2	3	4	5
	0	-1.0025	-1.1333	-0.9720	-0.7993	-0.6340	-0.4865
	1	-1.0014	-1.0468	-0.9087	-0.7593	-0.6148	-0.4844
	2	-1.0004	-0.9606	-0.8436	-0.7154	-0.5902	-0.4761
	3	-0.9994	-0.8754	-0.7769	-0.6682	-0.5609	-0.4621
	4	-0.9983	-0.7917	-0.7098	-0.6186	-0.5277	-0.4432
	5	-0.9973	-0.7104	-0.6431	-0.5674	-0.4913	-0.4199
	6	-0.9959	-0.6322	-0.5769	-0.5153	-0.4532	-0.3925
	7	-0.9948	-0.5567	-0.5132	-0.4636	-0.4120	-0.3645
	8	-0.9935	-0.4864	-0.4520	-0.4121	-0.3713	-0.3330
	9	-0.9921	-0.4200	-0.3941	-0.3635	-0.3312	-0.2996
	10	-0.9905	-0.3596	-0.3385	-0.3173	-0.2918	-0.2672
(b)	v	L					
		0	1	2	3	4	5
	0	0.0084	0.0046	-0.0010	-0.0070	-0.0129	-0.0183
	1	0.0057	0.0019	-0.0037	-0.0099	-0.0160	-0.0216
	2	0.0019	-0.0019	-0.0075	-0.0138	-0.0201	-0.0261
	3	-0.0034	-0.0071	-0.0128	-0.0192	-0.0257	-0.0319
	4	-0.0106	-0.0143	-0.0200	-0.0265	-0.0332	-0.0397
	5	-0.0203	-0.024	-0.0297	-0.0364	-0.0433	-0.0502
	6	-0.0353	-0.0372	-0.0431	-0.0498	-0.0570	-0.0644
	7	-0.0531	-0.0554	-0.0612	-0.0682	-0.0758	-0.0834
	8	-0.0780	-0.0802	-0.0863	-0.0937	-0.1017	-0.1100
	9	-0.1123	-0.1151	-0.1213	-0.1290	-0.1379	-0.1475
	10	-0.1609	-0.1645	-0.1715	-0.1795	-0.1897	-0.2009
(c)	v	L					
		0	1	2	3	4	5
	0	-0.12	-0.13	-0.15	-0.17	-0.19	-0.22
	1	-0.18	-0.18	-0.20	-0.22	-0.25	-0.28
	2	-0.24	-0.25	-0.27	-0.29	-0.31	-0.35
	3	-0.33	-0.34	-0.35	-0.38	-0.4	-0.44
	4	-0.44	-0.45	-0.47	-0.49	-0.52	-0.55
	5	-0.59	-0.60	-0.62	-0.64	-0.67	-0.71
	6	-0.81	-0.79	-0.81	-0.84	-0.87	-0.92
	7	-1.07	-1.05	-1.08	-1.11	-1.15	-1.19
	8	-1.42	-1.41	-1.43	-1.47	-1.52	-1.57
	9	-1.89	-1.89	-1.92	-1.97	-2.02	-2.10
	10	-2.56	-2.58	-2.62	-2.67	-2.74	-2.84
(d)	v	L					
		0	1	2	3	4	5
	0	-1.11	-2.14	11.27	1.79	1.12	0.91
	1	-2.32	-7.23	3.99	1.66	1.15	0.95
	2	-9.61	9.94	2.64	1.56	1.17	0.99
	3	7.39	3.55	2.07	1.47	1.18	1.02
	4	3.16	2.37	1.75	1.39	1.17	1.05
	5	2.18	1.87	1.55	1.32	1.16	1.06
	6	1.72	1.60	1.42	1.26	1.15	1.07
	7	1.50	1.43	1.32	1.22	1.13	1.07
	8	1.36	1.32	1.24	1.18	1.12	1.07
	9	1.27	1.23	1.19	1.14	1.10	1.07
	10	1.20	1.18	1.15	1.11	1.09	1.06

TABLE X: (a) Values of the dynamic vibrational contribution to the BBR shift of a level (v, L) of HD^+ , normalized to the static contribution, $\Delta E_{\text{dyn},rv}((v, L), T)/\Delta E_{\text{stat}}((v, L), T)$, at $T = 300$ K. (b) Approximate total BBR shift $\Delta E_{\text{stat}}((v, L), T) + \Delta E_{\text{dyn},rv}((v, L), T)$, in Hz. (c) temperature derivatives of the approximate total BBR shift, $d(\Delta E_{\text{stat}}((v, L), T) + \Delta E_{\text{dyn},rv}((v, L), T))/dT$, at 300 K, in mHz/K; (d) the normalized temperature derivative of the total BBR shift, $(T/4) \times (\Delta E_{\text{stat}} + \Delta E_{\text{dyn},rv})^{-1} \times d(\Delta E_{\text{stat}} + \Delta E_{\text{dyn},rv})/dT$, at 300 K. These results are non-relativistic and do not include the frequency-dependent contributions $\Delta E_{\text{dyn},elec}$ from excited electronic states, which are given in Tab. XI.

range $\omega \leq 0.08$.

We show in Fig. 2 the frequency-dependent part of the polarisability of one level of H_2^+ , $\alpha_{s,var,\text{H}_2^+}((1, 1), \omega) - \alpha_{s,var,\text{H}_2^+}((1, 1), 0)$, at low frequencies. For the computation of the BBR shift at 300 K, frequencies up to approximately $\omega = 0.013$ atomic units are relevant. In this range the polarisability is quite close to quadratic in ω . With increasing vibrational quantum numbers v , the deviations from quadratic are more pronounced.




FIG. 2: . Black (dotted line): the frequency-dependent contribution of the scalar polarisability of H_2^+ in the level $(v = 1, L = 1)$, computed using variational wave functions. Red (gray, full line): difference between the scalar polarisability of HD^+ , also in the level $(v = 1, L = 1)$, computed variationally and approximately. The difference is noticeable close to the two rotational transition frequencies $(v = 1, L = 1) \rightarrow (v = 1, L = 0)$ and $(v = 1, L = 1) \rightarrow (v = 1, L = 2)$. A gap occurs in the red curve at $\omega = 3.5 \times 10^{-4}$ atomic units because the difference is negative.

v	L	$\omega_{min}(v, L)$	fit	$\Delta E_{dyn,elec}((v, L), T)$ [mHz]
0	0	0.0015	$16.10 \omega^3 + 14.54 \omega^2$	-0.0021
0	1	0.0015	$16.24 \omega^3 + 14.62 \omega^2$	-0.0021
0	3	0.002	$16.94 \omega^3 + 15.01 \omega^2$	-0.0022
0	4	0.003	$17.58 \omega^3 + 15.32 \omega^2$	-0.0022
1	1	0.0015	$48.34 \omega^3 + 24.76 \omega^2$	-0.0036
1	5	0.019	$63.87 \omega^3 + 25.86 \omega^2$	-0.0038
2	4	0.02	$148.1 \omega^3 + 42.73 \omega^2$	-0.0063
2	5	0.019	$166.0 \omega^3 + 42.92 \omega^2$	-0.0064
3	2	0.04	$395.8 \omega^3 + 62.68 \omega^2$	-0.0094

TABLE XI: Fourth column: polynomial approximation to $\alpha_{s,dyn,elec}(\omega)$, the frequency-dependent part of the contribution to the polarisability α_s of HD^+ stemming from the excited electronic states. Fifth column: corresponding contribution to the BBR shift at 300 K. The angular frequencies ω and ω_{min} are in atomic units.

The dynamic electronic BBR shift corrections $\Delta E_{dyn,elec}$ computed from the variational data (with an integration analogous to Eq. (27)) are shown in Table VIII. We see that the correction is small in relative terms, 1×10^{-4} for $v = 0$, increasing to 4×10^{-4} for $v = 3$. It is very weakly dependent on L . Nevertheless, these results show that the dynamic contribution should not be omitted even within the non-relativistic approximation. When it is included, the overall inaccuracy is limited by the non-relativistic approximation to approximately 1×10^{-4} fractionally. For $v < 6$, $L < 6$ the total BBR shift is smaller than 0.1 Hz. Therefore, the absolute error is less than 0.01 mHz.

2. HD^+

For HD^+ , a comparison of the variational dynamic polarisability $\alpha_{s,var,\text{HD}^+}(\omega)$ with the approximation $\alpha_{s,\text{HD}^+}(\omega) = \alpha_s(\omega = 0) + \delta\alpha_{s,dyn,rv}(\omega)$ is depicted in Fig. 2, which shows their difference. In evaluating the approximation, we have used both the transition dipoles of Tian et al. [35] and their non-relativistic energies, since also the variational polarisability was computed in non-relativistic approximation. The agreement is very good, except for small deviations near the transition frequencies (whose nominal contribution to the BBR shift is only of order 1.5×10^{-3} mHz), and a frequency-dependent contribution from the excited electronic states, which is again closely quadratic in frequency.

We have fitted a simple quadratic plus cubic polynomial to the difference $\alpha_{s,var}(\omega) - \alpha_{s,\text{HD}^+}(\omega)$ between variational and approximate frequency-dependent polarisability, over the frequency range $\omega_{min}(v, L)$ to 0.05 at. u. Here, $\omega_{min}(v, L)$ is chosen appropriately so as to allow an accurate fit. These fits represent an approximation to $\delta\alpha_{s,dyn,elec}((v, L), \omega)$ for frequencies from 0 to 0.05 at. u. The fits are shown in Tab. XI. The contribution of the cubic term is seen to be small compared to the quadratic one for the range of frequencies relevant for the BBR shift

at 300 K. Tab. XI gives the corresponding contributions to the BBR shift, to be added to the other two contributions given in Tab. X. The error in the values of $\Delta E_{\text{dyn,elec}}$ due to this fit treatment is on the order of 0.001 mHz. We see that this BBR shift contribution again varies weakly with L , but significantly with v and that for levels with $v = 3$ it reaches 1×10^{-5} Hz. Therefore, it needs to be taken into account even within the non-relativistic approximation, if no loss of accuracy is desired. When this is done, the total error of the BBR shift due to the non-relativistic approximation is expected to be 1×10^{-4} fractionally, or less than 0.03 mHz for the low-lying levels of HD^+ , $v < 6$.

VI. CONCLUSION

We have computed the non-adiabatic static polarisabilities of the molecular hydrogen ions HD^+ , H_2^+ , and D_2^+ , extending significantly previous results, mostly limited to rovibrational levels with rotational angular momentum $L = 0, 1$. For a number of rovibrational levels, we have also computed the frequency-dependent non-adiabatic polarisability.

The dependence of the polarisabilities on the hyperfine state has been derived and discussed in detail. We have pointed out the special case of the pure states, for which a simple analytical result has been derived. This result is actually a very good approximation for all hyperfine states of H_2^+ . The hyperfine-state-dependence is of crucial importance if a detailed understanding of the systematic shifts of transition frequencies is to be performed.

We have also computed the shifts induced by the black-body radiation field, and their temperature derivatives.

Emphasis has been given here to achieve high numerical accuracy. The effective relative inaccuracy of our computed values is about 1×10^{-4} due to the neglect of relativistic corrections. For H_2^+ and D_2^+ this translates in an absolute inaccuracy of 0.001 at. u. for all levels with $v < 6$, $L < 5$. For HD^+ in $L = 0, 1$ levels the inaccuracy is less than 0.1 at. u., and in $L \geq 2$, it is less than 0.003 at. u.. An inaccuracy of 0.1 atomic unit is sufficiently low to allow evaluating the Stark shift with a theoretical error corresponding to the 10^{-18} fractional frequency level, given the typical electric field values in ion traps.

In order to obtain accurate values of the black-body radiation shift, we have used accurate values of the transition dipoles and we have analyzed the importance of the contributions from excited electronic states. We estimate the inaccuracy of the shifts to be less than 0.03 mHz for levels with $v < 6$, $L < 6$, at 300 K, for both HD^+ and H_2^+ . This corresponds to theoretical fractional frequency errors on the order of 1×10^{-18} .

Using the present results it becomes possible to identify theoretically transitions having low sensitivity to external fields [16, 17]. This represents an important aspect in the future spectroscopy of the simplest stable molecules.

-
- [1] B. Grémaud, D. Delande, and N. Billy, J. Phys. B **31**, 383 (1998).
 - [2] see V.I. Korobov, L. Hilico, and J.-Ph. Karr, “ $m\alpha^7$ -order QED corrections in the hydrogen molecular ions and antiprotonic helium”, Phys. Rev. Lett. **112**, 103003 (2014) and references therein.
 - [3] J.C.J. Koelemeij, B. Roth, A. Wicht, I. Ernsting, and S. Schiller, Phys. Rev. Lett. **98**, 173002 (2007).
 - [4] U. Fröhlich, B. Roth, P. Antonini, C. Lämmerzahl, A. Wicht, S. Schiller, in *Seminar on Astrophysics, Clocks and Fundamental Constants*, E. Peik, S. Karshenboim, eds., Lecture Notes in Physics, Springer, **648**, 297 (2004)
 - [5] S. Schiller and V.I. Korobov, Phys. Rev. A **71**, 032505 (2005).
 - [6] K. B. Jefferts, Phys. Rev. Lett. **23**, 1476 1969.
 - [7] W.H. Wing, G.A. Ruff, W.E. Lamb, Jr., and J. J. Spezeski, “Observation of the Infrared Spectrum of the Hydrogen Molecular Ion HD^+ ”, Phys. Rev. Lett. **36**, 1488–1491 (1976); doi/10.1103/PhysRevLett.36.1488;
 - [8] A. Carrington *et al.*, Mol. Phys. **72**, 735 (1991); doi: 10.1080/00268979100100531
 - [9] S. Schiller, C. Lämmerzahl, Phys. Rev. A **68**, 053406 (2003).
 - [10] P. Blythe, B. Roth, U. Fröhlich, H. Wenz, S. Schiller, “Production of cold trapped molecular hydrogen ions”, Phys. Rev. Lett. **95**, 183002 (2005); doi:10.1103/PhysRevLett.95.183002
 - [11] U. Bressel, A. Borodin, J. Shen, M. Hansen, I. Ernsting, and S. Schiller, “Manipulation of Individual Hyperfine States in Cold Trapped Molecular Ions and Application to HD^+ Frequency Metrology”, Phys. Rev. Lett. **108**, 183003 (2012)
 - [12] J. Shen, A. Borodin, M. Hansen, und S. Schiller, “Observation of a rotational transition of trapped and sympathetically cooled molecular ions”, Phys. Rev. A **85**, 032519 (2012).
 - [13] D. Bakalov, V.I. Korobov and S. Schiller, “Precision spectroscopy of the molecular ion HD^+ : control of Zeeman shifts”, Phys. Rev. A **82**, 055401 (2010).
 - [14] D. Bakalov, V.I. Korobov and S. Schiller, “Magnetic field effects in the transitions of the HD^+ molecular ion and precision spectroscopy”, J. Phys. B: At. Mol. Opt. Phys. **44**, 025003 (2011); Corrigendum: J. Phys. B: At. Mol. Opt. Phys. **45**, 049501 (2012).
 - [15] J.-P. Karr, V.I. Korobov, L. Hilico, “Vibrational spectroscopy of H_2^+ : Precise evaluation of the Zeeman effect”, Phys. Rev. A **77**, 062507 (2008)

- [16] D. Bakalov and S. Schiller, “The electric quadrupole moment of molecular hydrogen ions and their potential for a molecular ion clock”, Appl. Phys. B 114, 213-230 (2014); DOI 10.1007/s00340-013-5703-z
- [17] S. Schiller, D. Bakalov, V. I. Korobov, arXiv:1402.1789; Phys. Rev. Lett. (accepted).
- [18] J.C.J. Koelemeij, “Infrared dynamic polarisability of HD^+ rovibrational states”, Phys. Chem. Chem. Phys. **13**, 18844 (2011).
- [19] D. M. Bishop and B. Lam, “An analysis of the interaction between a distant point charge and H_2^+ ”, Mol. Phys. 65, 679-688 (1988)
- [20] L. Hilico, N. Billy, B. Grémaud, D. Delande, “Polarisabilities, light shifts and two-photon transition probabilities between $J = 0$ states of the H_2^+ and D_2^+ molecular ions”, J. Phys. B. 34, 491 (2001)
- [21] R.E. Moss, and L. Valenzano, “The dipole polarisability of the hydrogen molecular cation HD^+ and other isotopomers”, Molec. Phys. **100**, 1527 (2002).
- [22] Z.C. Yan, Y.J. Zhang, and Y. Li, Phys. Rev. A 67, 062504 (2003)
- [23] J.-Ph. Karr, S. Kilic, and L. Hilico, “Energy levels and two-photon transition probabilities in the HD^+ ion”, J. Phys. B: At. Mol. Opt. Phys. **38**, 853 (2005).
- [24] A. K. Bhatia, R. J. Drachman, Phys. Rev. A 61, 032503 (2000)
- [25] V.I. Korobov, Phys. Rev. A **61**, 064503 (2000)
- [26] D. Bakalov and S. Schiller, “Static Stark effect in the molecular ion HD^+ ”, Hyperfine interact. **210**, 25 (2012)
- [27] H. Olivares Pilón and D. Baye, “Static and dynamic polarisabilities of the non-relativistic hydrogen molecular ion”, J. Phys. B: At. Mol. Opt. Phys. 45, 235101 (2012); <http://iopscience.iop.org/0953-4075/45/23/235101>
- [28] This result was first presented at the 2nd COST IOTA Workshop on Cold Molecular Ions, Arosa (Switzerland), September 2013.
- [29] V.I. Korobov, L. Hilico, and J.-Ph. Karr, Phys. Rev. A **74**, 040502(R) (2006).
- [30] J. Ph. Karr, F. Bielsa, A. Douillet, J. Pedregosa, V. I. Korobov, and L. Hilico, Phys. Rev. A 77, 063410 (2008).
- [31] D. Bakalov, V.I. Korobov, and S. Schiller, “High-precision calculation of the hyperfine structure of the HD^+ ion”, Phys. Rev. Lett. 97, 243001 (2006).
- [32] A Messiah, *Quantum Mechanics*, (vol. II, Appendix C, § 15), North Holland, Amsterdam, 1961.
- [33] D.A. Varshalovich, A.N. Moskalev, and V.K. Khersonskii, *Quantum Theory of Angular Momentum*, (World Scientific, Singapore, 1988).
- [34] I. Roeggen, “The theory of the Stark effect in multiplet states of diatomic polar molecules”, J. Phys. B: At. Mol. Opt. Phys. 4, 168 (1971); <http://iopscience.iop.org/0022-3700/4/2/004>
- [35] Q.-L. Tian, L.-Y. Tang, Z.-X. Zhong, Z.-C. Yan, T.-Y. Shi, “Oscillator strengths between low-lying ro-vibrational states of hydrogen molecular ions”. J. of Chem. Phys. 137, 024311 (2012). doi:10.1063/1.4733988.
- [36] V.I. Korobov, “Leading-order relativistic corrections to the ro-vibrational spectrum of H_2^+ and HD^+ molecular ions”, Phys. Rev. A 74, 052506 (2006).
- [37] V.I. Korobov and Zhen-Xiang Zhong, “Bethe Logarithm for the H_2^+ and HD^+ molecular ions, Phys. Rev. A 86, 044501 (2012).
- [38] R. E. Moss, “Calculations for vibration-rotation levels of HD^+ , in particular for high N“. Molecular Physics 78, 371–405. doi:10.1080/00268979300100291.
- [39] T.N. Rescigno, and V. McKoy, Phys. Rev. A 12, 522 (1975).
- [40] V.I. Korobov, “Dynamic polarisability properties of the weakly bound ddu and dtu molecular ions”, J. Phys. B 37, 2331–2341 (2004), DOI: 10.1088/0953-4075/37/11/010
- [41] Online at: (**** to be filled by editor ****). Each line of the data files has the format: ω (in atomic units), $-\alpha_s((v, L), \omega)$ (in atomic units), $-\alpha_t((v, L), \omega)$ (in atomic units). Note that for HD^+ and $v = 3, L = 2$ there is a difference of 0.0001 at. u. between the $\omega = 0$ value of the data file and the value in I. This arises from the numerical procedure used, which was optimized towards good accuracy for high ω up to the continuum threshold. Note that the values of α_s, α_t in the files should be multiplied with (-1) for actual computations.
- [42] H.O. Pilón, priv. comm. (2013)